



022-CHM

## Quantum chemical study of Pnicogen bond based on point-of-charge and MEP

Mahmoud A. A. Ibrahim, Hussein M. H. Ali, Abeer A. M. Hasb  
Chemistry Department, Faculty of Science, Minia University, Minia 61519, Egypt

*A pnicogen bond is a noncovalent bond between a pnicogen atom (X) and a Lewis base (Y). This type of bond is attributed to the anisotropic distribution of the charge density on the pnicogen atom, resulting in the formation of a positive cap (called the  $\sigma$ -hole)[1]. Pnicogen bond is similar to halogen and chalcogen[2] bond in that they contribute to the stabilization of macromolecules and inhibitor-receptor complexes. Pnicogen bond's characteristics are still unrevealed. For instance, the ability of PH<sub>3</sub> molecule to form a dimer via pnicogen bond was under debate. In current work, six molecules (XH<sub>3</sub>, XH<sub>2</sub>F and XF<sub>3</sub>; X = N and P) were studied using quantum mechanical calculations including point-of-charge (PoC), stabilization energy, natural bonding orbital (NBO) and Molecular electrostatic potential (MEP) calculations. Firstly, all molecules were fully optimized by the second-order Møller–Plesset (MP2) method with 6-311++G\*\* basis set. MEPs were then generated for the molecules at B3LYP/6-311++G\*\* level of theory. The stabilization energy of the studied molecules was calculated at MP2/aug-cc-pVTZ level with incorporation of a PoC at a distance ranging from 1.5 to 8.0 Å. The results showed that the existence of a negative charge, i.e. Lewis base, causes destabilization in case of NH<sub>3</sub> and PH<sub>3</sub> molecules. In case of NH<sub>2</sub>F, NF<sub>3</sub>, PH<sub>2</sub>F and PF<sub>3</sub>, a PoC with a value of -0.25au enhances the stabilization of molecular energy with a value of ranging from -1.0 to -2.6 kcal/mol. Furthermore, quantum calculations including MEP and NBO showed that NH<sub>3</sub> and PH<sub>3</sub> molecules are not able to form pnicogen bond. The impact of Y-X...PoC angle and value of PoC on the stabilization energy was evaluated. The current study revealed the main pnicogen bond's characteristics which would serve as a framework for all future studies.*